

Autism, a variant of kernicterus?

Eileen Nicole Simon, Ph.D., R.N.,

Stereotyped motor mannerisms in children with autism may reflect impairment of the basal ganglia (Langen et al., 2007; Rinehart et al., 2006; Hollander et al., 2005).

Damage of the basal ganglia, associated with jaundice in the newborn period, leads to kernicterus, a severe affliction of motor development (Shapiro, 2005). Hemolytic disease of the newborn, caused by maternal antibodies that cross the placenta, is the best known cause of kernicterus (Eder, 2006). However, development of RhoGAM to prevent maternal antibody formation to Rh-factor has not eradicated the problem (Ip et al., 2004). Hansen (2000) pointed to an increase in kernicterus since 1994, which he even referred to as a new "epidemic" (p1156).

Jaundice caused by elevated bilirubin levels is common in newborns. Why damage occurs in some but not all infants remains an important question (Maisels, 2006). Zimmerman and Yannet (1933) summarized a large number of case reports and concluded that injury by anoxia or sepsis often precedes bilirubin staining of subcortical nuclei, and stated, "This differs in no way from the well known fact that any intravital dye will localize in zones of injury and will leave unstained tissues which are not damaged," (p757).

The blood-brain barrier normally prevents bilirubin from getting into the brain anywhere, but if breached, according to Levine et al. (1982), "Other toxic substances may also enter the brain, but they could go unnoticed if they are colorless." (p258). Thus substances other than bilirubin should be looked for that could result in kernicterus-like injury of subcortical nuclei.

Kernicterus has been reported in infants with very low levels of bilirubin (Harris, Lucey, & MacLean, 1958), and kernicterus was found in newborn infants following treatment with sulfisoxazole antibiotic, and with synthetic vitamin K (Robertson, 2003a, b, c). Factors that might trigger breakdown of the blood-brain barrier should be looked for as the real cause of kernicterus.

The basal ganglia are among a rank order of subcortical structures that comprise the most metabolically active sites in the brain (Sokoloff, 1981; Kennedy, Sakurada, Shinohara, & Miyaoka, 1982). Factors that disrupt aerobic metabolism during development also impair these centers of high metabolic rate (Kornreich et al., 2005; Clark et al., 2000). The same subcortical nuclei were also damaged in experiments with monkeys on asphyxia at birth, in the first of which Ranck and Windle (1959) noted, "The human neuropathologic entity most closely resembling the effects of asphyxia neonatorum in the monkey is kernicterus" (p153).

Lucey et al. (1964) injected monkeys with bilirubin "every 6 hours from 1 to 3 hours after birth for as long as 4 days" (p43), and neither kernicterus nor any neurological signs were seen. Monkeys subjected to asphyxia before being made hyperbilirubinemic, developed severe neurological deficits and were found to have kernicterus.

Evidence of anoxic birth in children who later develop autism is abundant (Badawi et al., 2006; Larsson et al., 2005; Glasson et al., 2004). Could the repetitive and stereotyped mannerisms observed in children with autism represent a variant form of kernicterus?

References

- Badawi, N., Dixon, G., Felix, J.F., Keogh, J.M., Petterson, B., Stanley, F.J., & Kurinczuk, J.J. (2006). Autism following a history of newborn encephalopathy: more than a coincidence? *Developmental Medicine and Child Neurology*, *48*, 85-89.
- Clark, C.M., Li, D., Conry, J., Conry, R., & Looock, C. (2000). Structural and functional brain integrity of fetal alcohol syndrome in nonretarded cases. *Pediatrics*, *105*, 1096-1069.
- Eder, A.F. (2006). Update on HDFN: new information on long-standing controversies. *Immunohematology /American Red Cross*, *22*, 188-195.
- Glasson, E. J., Bower, C., Petterson, B., de Klerk, N., Chaney, G., & Hallmayer, J.F. (2004). Perinatal factors and the development of autism: a population study. *Archives of General Psychiatry*, *61*, 618-627.
- Hansen, T.W. (2000) Kernicterus in term and near-term infants--the specter walks again. *Acta Paediatrica*, *89*, 1155-1157.
- Harris, R.C., Lucey, J.F., & MacLean, J.R. (1958). Kernicterus in premature infants associated with low concentrations of bilirubin in the plasma. *Pediatrics*, *21*, 875-884.
- Hollander, E., Anagnostou, E., Chaplin, W., Esposito, K., Haznedar, M.M., Licalzi, E., Wasserman, S., Soorya, L., & Buchsbaum M. (2005). Striatal volume on magnetic resonance imaging and repetitive behaviors in autism. *Biological Psychiatry*, *58*, 226-232.
- Ip, S., Lau, J., Chung, M., Kulig, J., Sege, R., Glicker, S., & O'Brien, R. (2004) Hyperbilirubinemia and kernicterus: 50 years later. *Pediatrics*, *114*,263-264.
- Kennedy, C., Sakurada, O., Shinohara, M., & Miyaoka, M. (1982) Local cerebral glucose utilization in the newborn macaque monkey. *Annals of Neurology*, *12*, 333-340.
- Kornreich, L., Bron-Harlev, E., Hoffmann, C., Schwarz, M., Konen, O., Schoenfeld, T., Straussberg, R., Nahum, E., Ibrahim, A.K., Eshel, G., & Horev, G. (2005) Thiamine deficiency in infants: MR findings in the brain. *AJNR American Journal of Neuroradiology*, *26*, 1668-1674.
- Langen, M., Durston, S., Staal, W.G., Palmen, S.J., van Engeland, H. (2007). Caudate nucleus is enlarged in high-functioning medication-naive subjects with autism. *Biological Psychiatry*, *62*, 262-266.
- Larsson, H.J., Eaton, W.W., Madsen, K.M., Vestergaard, M., Olesen, A.V., Agerbo, E., Schendel, D., Thorsen, P., & Mortensen, P.B. (2005). Risk factors for autism: perinatal factors, parental psychiatric history, and socioeconomic status. *American Journal of Epidemiology*. *161*, 916-925.
- Levine, R.L., Fredericks, W.R., & Rapoport, S. (1982) Entry of bilirubin into brain due to opening of the blood-brain barrier. *Pediatrics*, *69*, 255-259.
- Lucey, J.F., Hibbard, E., Behrman, R.E., Esquivel de Gallardo, F.O., & Windle, W.F. (1964). Kernicterus in asphyxiated newborn rhesus monkeys. *Experimental Neurology* *9*, 43-58.
- Maisels, M.J. (2006). What's in a name? Physiologic and pathologic jaundice: the conundrum of defining normal bilirubin levels in the newborn. *Pediatrics*, *118*, 805-807.

- Ranck, J.B. Jr. & Windle, W.F. (1959). Brain damage in the monkey, macaca mulatta, by asphyxia neonatorum. *Experimental Neurology* 1,130-154.
- Rinehart, N.J., Tonge, B.J., Ianssek, R., McGinley, J., Brereton, A.V., Enticott, P.G., & Bradshaw, J.L. (2006). Gait function in newly diagnosed children with autism: Cerebellar and basal ganglia related motor disorder. *Developmental Medicine and Child Neurology*, 48, 819-824.
- Robertson, A.F. (2003a) Reflections on errors in neonatology: I. The "Hands-Off" years, 1920 to 1950. *Journal of Perinatology*, 23, 48-55.
- Robertson, A.F. (2003b).Reflections on errors in neonatology: II. The "Heroic" years, 1950 to 1970. *Journal of Perinatology*, 23, 154-161.
- Robertson, A.F. (2003c).Reflections on errors in neonatology III. The "experienced" years, 1970 to 2000. *Journal of Perinatology*, 23, 240-249.
- Shapiro, S.M. (2005). Definition of the clinical spectrum of kernicterus and bilirubin-induced neurologic dysfunction (BIND). *Journal of Perinatology*, 25, 54-59.
- Sokoloff, L. (1981). Localization of functional activity in the central nervous system by measurement of glucose utilization with radioactive deoxyglucose. *Journal of Cerebral Blood Flow and Metabolism*, 1, 7-36.